

略語表

| | |
|---------------|--|
| ADS | Agitation Distress Scale |
| AUC | area under the curve |
| BCS | Bedside Confusion Scale |
| CAM | Confusion Assessment Method |
| CAM-ICU | the Confusion Assessment Method for the Intensive Care Unit |
| CCS | Communication Capacity Scale |
| CDT | Clock Drawing Test |
| CGI-S | Clinical Global Impression-Severity |
| CRS | Comprehensive Ranking System |
| CTCAE | Common Terminology Criteria for Adverse Events |
| CTD | Cognitive Test for Delirium |
| DELTA | Delirium Team Approach |
| DOS | Delirium Observation Screening Scale |
| DMS | Delirium Motoric Checklist |
| DRS-c | Delirium Rating Scale-Chinese |
| DRS-R-98 | Delirium Rating Scale-Revised-98 |
| DSM | Diagnostic and Statistical Manual of Mental Disorders |
| ECOG | Eastern Cooperative Oncology Group |
| EORTC QLQ-C30 | European Organization for Research and Treatment of Cancer Quality of Life Questionnaire-Core 30 |
| GHQ-28 | General Health Questionnaire-28 |
| HADS | Hospital Anxiety and Depression Scale |
| ICD | International Statistical Classification of Diseases and Related Health Problems |
| MAC | Mental Adjustment to Cancer |
| MARTA | multi-acting receptor-targeted antipsychotics |
| MDAS | Memorial Delirium Assessment Scale |
| MMSE | Mini-Mental State Examination |
| NAS | Numeric Analog Scale |
| NSAIDs | non-steroidal anti-inflammatory drugs |
| Nu-DESC | Nursing Delirium Screening Scale |
| RASS | Richmond Agitation-Sedation Scale |

(つづく)

| | |
|--------|---|
| RBANS | Repeatable Battery for the Assessment of Neuropsychological Status |
| ROC | receiver operating characteristic |
| STAS-J | Japanese version of the Support Team Assessment Schedule |
| SQiD | Single Question in Delirium |
| VAS | Visual Analog Scale |
| 3D-CAM | the 3-Minute Diagnostic Assessment for Delirium using the CAM algorithm |